

Osteopontin-Induced, Integrin-Dependent Migration of Human Mammary Epithelial Cells Involves Activation of the Hepatocyte Growth Factor Receptor (Met)

Alan B. Tuck,^{1,2,3,4*} Bruce E. Elliott,⁵ Charulata Hota,^{1,2,3,4} Eric Tremblay,⁵ and Ann F. Chambers^{1,2,4}

¹Department of Pathology, University of Western Ontario, London, Ontario, Canada N6A 5C1

²Department of Oncology, University of Western Ontario, London, Ontario, Canada N6A 5C1

³London Health Sciences Centre, London, Ontario, Canada N6A 4G5

⁴London Regional Cancer Centre, London, Ontario, Canada N6A 4L6

⁵Cancer Research Labs, Queen's University, Kingston, Ontario, Canada K7L 3N6

Abstract Osteopontin (OPN) is a secreted glycoposphoprotein which induces migration of mammary carcinoma cells, and has been implicated in the malignancy of breast carcinoma. Hepatocyte growth factor (HGF) induces cell migration of several mammary epithelial cell (MEC) lines, via activation of its cognate receptor (Met). This study examines the mechanism of OPN-induced MEC migration, in terms of the cell surface integrins involved and induction of the HGF/Met pathway. Three different MEC cell lines were used, representing different stages of tumor progression: 21PT, non-tumorigenic; 21NT, tumorigenic; non-metastatic; and MDA-MB-435, tumorigenic, highly metastatic. Human recombinant OPN was found to induce the migration of all three lines. OPN-induced migration of 21PT and 21NT cells was $\alpha\beta 5$ and $\beta 1$ -integrin dependent, and $\alpha\beta 3$ -independent, while that of MDA-MB-435 cells was $\alpha\beta 3$ -dependent. HGF also induced migration of all three cell lines, and a synergistic response was seen to HGF and OPN together. The increased migration response to OPN was found to be associated with an initial increase in Met kinase activity (within 30 min), followed by an increase in Met mRNA and protein expression. OPN-induced cell migration is thus mediated by different cell surface integrins in MEC lines representing different stages of progression, and involves activation of the HGF receptor, Met. *J. Cell. Biochem.* 78:465–475, 2000. © 2000 Wiley-Liss, Inc.

Key words: osteopontin (OPN); cell migration; integrin; hepatocyte growth factor (HGF); Met; mammary epithelial cells; breast cancer

Growth, migration, and differentiation of epithelial cells are known to be dependent upon integrin-mediated adhesion to extracellular matrix components [reviews in Assoian, 1997; Bissell, 1999; Gumbiner, 1996]. Similarly, these same cellular processes are also known to

be influenced by a number of different growth factor pathways [reviews in Sedorf, 1995; Vande Woude et al., 1997; Heldin, 1998; Birchmeier, 1998]. Several recent studies have addressed the possibility of interactions between integrin and growth factor mediated pathways, with evidence emerging for both growth factor control of cell adhesion events [van der Voort et al., 1997; Weimar et al., 1997; Trusolino et al., 1998; Weimar et al., 1998], and conversely, for integrin-mediated cell adhesion phenomena influencing sensitivity to certain growth factors [Miyamoto et al., 1996; Brooks et al., 1997]. However, the mechanism and biological relevance of these growth factor-integrin interactions are not yet clear.

Our group has particular interest in the role of the secreted glycoposphoprotein OPN in the malignancy of breast cancer. We have found that OPN can induce cell migration and

Grant sponsor: Canadian Breast Cancer Research Initiative; Grant number: 8426; Grant sponsor: U.S. Army Medical Research and Materials Command; Grant number: DAMD17-96-1-6251; Grant sponsor: U.S. Army Breast Cancer Research Program; Grant number: DAMD17-96-1-6076.

The content of this article does not necessarily reflect the position or policy of the U.S. government, and no official endorsement should be inferred.

*Correspondence to: Alan B. Tuck, MD, PhD, Department of Pathology, London Health Sciences Centre, Victoria Campus, 375 South St., London, Ontario N6A 4G5, Canada. E-mail: atuck@julian.uwo.ca

Received 6 December 1999; Accepted 20 March 2000

Print compilation © 2000 Wiley-Liss, Inc.

invasiveness of cultured mammary epithelial cells (MECs) [Xuan et al., 1994; Xuan et al., 1995; Tuck et al., 1999], that it may be secreted in greater quantity by MECs of greater degree of malignancy [Tuck et al., 1999], and that higher levels (tumor cell or plasma levels respectively) are associated with poorer prognosis in patients with either lymph node negative or metastatic breast cancer [Tuck et al., 1998; Singhal et al., 1997]. Evidence from us [Xuan et al., 1994, 1995; Tuck et al., 1999], and others [Senger et al., 1996], has indicated that OPN-induced cell movement is a directed, RGD-dependent response, although CD44-mediated phenomena may also be involved [Weber et al., 1996; Bourguignon et al., 1998, 1999; Katagiri et al., 1999; Tuck et al., unpublished observations]. Cell adhesion studies have shown for a variety of cell types that the major cell surface integrins involved in OPN binding include $\alpha v \beta 1$, $\alpha v \beta 3$, and $\alpha v \beta 5$ [Hu et al., 1995; Liaw et al., 1995].

Given this information, along with the abundant evidence for the importance of HGF/Met in cell motility of MECs [Bhargava et al., 1992; Rosen et al., 1994; Rahimi et al., 1998], and in the malignancy of breast cancer [Yamashita et al., 1994; Tuck et al., 1996; Yao et al., 1996; Jin et al., 1997; Bevilgia et al., 1997; Ghoussoub et al., 1998], we set out to examine the nature of OPN-induced cell migration, with respect to the involvement of cell surface integrins known to bind OPN, and possible interactions with the HGF/Met pathway. We have made use of three MEC lines, of differing malignancy: 21PT, non-tumorigenic; 21NT, tumorigenic, non-metastatic [Band et al., 1990]; and MDA-MB-435: tumorigenic, highly metastatic [Price et al., 1990]. We have assessed these cells for migratory responsiveness to OPN, alone and in combination with HGF. Having found evidence for a synergistic relationship between OPN and HGF in inducing cell migration, we proceeded to characterize the cell surface integrins involved, using blocking antibodies to $\alpha v \beta 5$, $\beta 1$, or $\alpha v \beta 3$ integrins. OPN-treated cells were then examined in time course experiments for induction of Met kinase activity and tyrosine phosphorylation, and for levels of HGF and Met mRNA and protein. Incubation with OPN was found to result in rapid activation of Met (all three cell lines), followed by an increase in Met RNA (all three cell lines) and protein (21PT and 21NT).

This work thus provides evidence that MEC cell lines representative of different stages of progression make use of different cell surface integrins in the migration response to OPN, and that this OPN-induced cell migration may be mediated at least in part by activation of Met.

METHODS

Cell Lines and Culture

The 21T series cell lines (21PT, 21NT) were obtained as a kind gift of Dr. Vimla Band (Dana Farber Cancer Institute) [Band et al., 1990]. These cells were maintained in culture in α -MEM supplemented with 10% FCS, 2 mM L-glutamine (all from GIBCO-BRL/Life Technologies, Grand Island, NY), insulin (1 μ g/ml), epidermal growth factor (EGF; 12.5 ng/ml), hydrocortisone (2.8 μ M), 10 mM HEPES, 1 mM sodium pyruvate, 0.1 mM nonessential amino acids, and 50 μ g/ml gentamycin (all from Sigma; α HE medium). MDA-MB-435 cells [Price et al., 1990] were obtained as a kind gift of Dr. Janet Price (MD Anderson Cancer Center, Houston, TX), and were grown in α -MEM with 10% FCS (both from GIBCO-BRL/Life Technologies).

Cell Migration

Cell migration assays were performed essentially as described previously [Xuan et al., 1995], using 24-well transwell chambers with polycarbonate filters of 8 μ m pore size (Costar, Cambridge, MA). Gelatin (Sigma) was applied at 6 μ g/filter and air dried. The gelatin was rehydrated with 100 μ l of serum-free α HE medium at room temperature for 90 min. Lower wells contained 800 μ l of α HE plus 0.1% BSA, with or without OPN, HGF, and/or blocking antibodies (as specified in Figs. 1 and 2). Human OPN (50 μ g/ml) used was the full length human recombinant GST-OPN (hrOPN), as previously described [Xuan et al., 1994]. Previous control experiments have shown that the GST portion alone has no influence on migration of these cells. Human HGF (20 ng/ml) was obtained from Collaborative Biomedical Products (Becton-Dickinson, Bedford, MA). Blocking anti-integrin antibodies included anti- $\alpha v \beta 3$ (Cedarlane, Hornby, ON), anti- $\alpha v \beta 5$ (GIBCO-BRL), and anti- $\beta 1$ (GIBCO-BRL), all used at saturating concentrations as determined by preliminary titration experiments. Cells

(5×10^4) were added to each upper well in α HE medium with 0.1% BSA and incubated for 5 h at 37°C. At the end of the incubation time, the cells that had migrated to the undersurface of the filters were fixed in place with glutaraldehyde and stained with hematoxylin. Cells that had not migrated and were attached to the

upper surface of the filters were removed from the filters with a cotton swab. The lower surfaces of the filters were examined microscopically under 100 \times magnification and representative areas were counted to determine the number of cells that had migrated through the filters. Control experiments were also performed in which blocking antibody in the lower chamber was replaced by non-immune mouse IgG (Cedarlane) at comparable concentration.

All cell migration and invasion assays were performed in triplicate. Statistical differences between groups were assessed using Student's *t*-test, with SigmaStat (Jandel Scientific, San Rafael, CA) statistical software.

Immunoprecipitation and Western Blotting for Met and Phosphotyrosine

Cells in monolayer were grown to 85–90% confluence, serum starved overnight, and incubated in serum-free medium either with or without human OPN (50 μ g/ml) or HGF (20 ng/ml) for the times specified. Cells were then rinsed with cold PBS, and lysed in lysis buffer containing 50 mM Tris-HCL (pH 7.5), 150 mM NaCl, 1% NP-40, 1 mM Na_3VO_4 , 50 mM NaF, 2 mM EGTA, 2 μ g/ml aprotinin, 2 μ g/ml leupeptin, and 1 mM PMSF. Lysates were centrifuged for 10 min at 14,000 rpm in an IEC/Micromax centrifuge at 4°C. Protein concentration of supernatants was determined using a bicinchoninic acid protein assay (Pierce, Rockford, IL). Equal protein amounts of each lysate were immunoprecipitated with rabbit anti-human Met polyclonal antibody at 4°C for 2 h.

Immunoprecipitates were collected on protein A-Sepharose (Amersham-Pharmacia Bio-




Fig. 1. a: OPN-induced migration of 21PT (PT) and 21NT (NT) cells is α v β 5, not α v β 3, integrin-dependent. Migration assays were performed as described in Materials and Methods. Lower chamber conditions were as follows: 0.1% BSA only (0); 50 μ g/ml hrOPN only (OP); 50 μ g/ml hrOPN with 15 μ g/ml non-specific mouse IgG (OP+nlG); 50 μ g/ml hrOPN with 30 μ g/ml anti- α v β 3 integrin blocking antibody (OP+ α v β 3); or 50 μ g/ml hrOPN with 15 μ g/ml anti- α v β 5 integrin blocking antibody (OP+ α v β 5). b: OPN-induced migration of 21PT (PT) and 21NT (NT) cells is β 1 integrin-dependent. Lower chamber conditions were as follows: 0.1% BSA only (0); 50 μ g/ml hrOPN only (OP); 50 μ g/ml hrOPN with 15 μ g/ml non-specific mouse IgG (OP+nlG); or 50 μ g/ml hrOPN with 15 μ g/ml anti- β 1 integrin blocking antibody (OP+ α β 1). Bar graphs represent the mean of four or five counts from each of three separate wells; error bars are SEM.




Fig. 2. OPN-induced migration of MDA-MB-435 (MDA) cells is $\alpha\beta 3$, not $\alpha\beta 5$ or $\beta 1$ integrin-dependent. Migration assays were performed as described in Materials and Methods. Lower chamber conditions were as follows: 0.1% BSA only (0); 50 $\mu\text{g/ml}$ hrOPN only (OP); 50 $\mu\text{g/ml}$ hrOPN with 25 $\mu\text{g/ml}$ non-specific mouse IgG (OP+nlG); 50 $\mu\text{g/ml}$ OPN with 15 $\mu\text{g/ml}$ anti- $\beta 1$ integrin blocking antibody (OP+ $\alpha\beta 1$); 50 $\mu\text{g/ml}$ hrOPN with 25 $\mu\text{g/ml}$ anti- $\alpha\beta 3$ integrin blocking antibody (OP+ $\alpha\alpha\beta 3$) or 50 $\mu\text{g/ml}$ hrOPN with 15 $\mu\text{g/ml}$ anti- $\alpha\beta 5$ integrin blocking antibody (OP+ $\alpha\alpha\beta 5$). Bar graphs represent the mean of four or five counts from each of three separate wells; error bars are SEM.

tech, Baie d'Urfe, Quebec, Canada), washed three times with lysis buffer, separated by 7% SDS-PAGE, and transferred to a nitrocellulose membrane. The membrane was blocked for 15 min with 3% skim milk, or 1% BSA, in TBST (10 mM Tris-HCl, pH 8.0, 150 mM NaCl, 0.1% Tween 20), and probed for 1 h with either mouse anti-human Met (DL-21 clone, Upstate Biotechnology Inc., Lake Placid, NY) or anti-phosphotyrosine antibody (PY20 clone, Transduction Labs, Lexington, KY). The membrane was washed three times for 5 min each with TBST buffer, incubated with horseradish peroxidase-labeled secondary anti-mouse antibody (Amersham-Pharmacia Biotech) for 15 min, and washed three times with TBST for

10 min each. Immune complexes were detected using ECL (Mandel/NEN, Guelph, ON).

In Vitro Met Kinase Assay

Cell cultures incubated in the presence or absence of hrOPN (50 $\mu\text{g/ml}$) or HGF (20 ng/ml) were rinsed with cold PBS, lysed, and immunoprecipitated as above. Immunoprecipitates were washed twice with cold lysis buffer and once with cold kinase buffer (20 mM PIPES, pH 7.0, 10 mM MnCl_2 , 10 μM Na_2VO_4). In vitro Met kinase activity was determined by incubating immunoprecipitates with 20 μl of kinase buffer containing 10 μCi [γ - ^{32}P] ATP at 30°C for 10 min. The reaction was stopped by addition of 2× SDS sample buffer containing 5% β -mercaptoethanol. Samples were boiled for 3 min and subjected to 7% SDS-PAGE. Serine and threonine phosphorylations were hydrolyzed by incubating the acrylamide gel in 1 M KOH at 45°C for 30 min, followed by fixing in 45% MeOH and 10% acetic acid for 30 min at room temperature and drying for 2 h at 80°C under a vacuum. Autoradiograms were produced and quantitated using a Storm Phosphorimager (Molecular Dynamics, Sunnyvale, CA).

Analysis of Met and HGF mRNA Levels

Cell cultures (85–90% confluent) were incubated in serum-free medium for the specified times in the presence or absence of 50 $\mu\text{g/ml}$ hrOPN. Cells were harvested by gentle trypsinization, pelleted, and mechanically homogenized (Polytron PT 1200, Brinkman Instruments [Canada] Ltd., Mississauga, ON). RNA was extracted using TRIzol Reagent (Canadian Life Technologies Inc., Burlington, ON), according to the protocol supplied by the manufacturer. RNA (10 $\mu\text{g/lane}$) was run on a 1.1% agarose gel with 6.8% formaldehyde, and capillary-transferred to GeneScreen Plus filters (DuPont Canada Inc., Mississauga, ON). Blots were probed with denatured, oligolabeled [α - ^{32}P]-dCTP cDNA probes (labeled using a kit provided by Pharmacia), according to the procedures provided by the manufacturers, and as previously described [Tuck et al., 1990, 1991]. cDNA probes were as follows: hepatocyte growth factor (HGF)—540 bp BamHI-XhoI fragment of human HGF cDNA [Nakamura et al., 1989]; Met/HGF receptor (HGFR)—800 bp EcoRI-EcoRV fragment of the human *met* cDNA [Park et al., 1987]; 18S rRNA (18S)—from p100D9.

Integrin-Dependent, OPN-Induced Migration Involves Met

469




Fig. 3. Synergistic effect of OPN and HGF on migration of 21PT (PT), 21NT (NT), and MDA-MB-435 (MDA) cells. Migration assays were performed as described in Materials and Methods. Contents of the lower chamber consisted of either: medium (aH, no EGF) without HGF or hrOPN (0); medium with 50 μ g/ml hrOPN (OP); medium with 10 ng/ml HGF (HGF); or medium with 50 μ g/ml hrOPN and 10 ng/ml HGF (OP+HGF). Bar graphs represent the mean of four or five counts from each of three separate wells; error bars are SEM.

RESULTS

OPN-Induced Migration of 21PT and 21NT Cells Involves Different Cell Surface Integrins Than for MDA-MB-435 Cells

Cell migration of 21PT, 21NT, and MDA-MB-435 cells was found to occur in response to hrOPN at a level comparable to that determined previously [Tuck et al., 1999]. Blocking experiments were performed using saturating concentrations of anti-integrin antibodies in the lower chamber of transwells, as described in Methods. For 21PT and 21NT cells, complete blocking of OPN-induced cell migration (to baseline levels) was obtained with the anti- α β 5 and β 1 integrin antibodies (Fig. 1a,b; $P < 0.002$ for all, Student's *t*-test). In contrast, non-immune mouse IgG did not block migration of either cell line. Saturating concentrations (30 μ g/ml) of anti- α β 3 integrin antibody had no detectable effect on migration of either 21PT or 21NT cells (Fig. 1a).




Fig. 4. HGF (a) and OPN (b)-induced increase in total cellular Met kinase activity of 21PT and 21NT cells. OPN (c)-induced increase in total Met protein of 21PT and 21NT cells. a,b: Cells were incubated ± 20 ng/ml HGF or 50 μ g/ml hrOPN for the times indicated, and cells lysates were prepared. Equal protein amounts of each lysate were immunoprecipitated with anti-Met IgG, and in vitro Met kinase activity was determined as described in Materials and Methods. Quantitation was done using a Phosphorimager. Total Met kinase activity is expressed in cpm/sample. c: Met protein was quantitated by immunoprecipitation with rabbit polyclonal anti-Met antibody, followed by 7% SDS-PAGE and Western blotting as described in Materials and Methods. Total Met protein was quantitated by densitometry and is expressed in integrated density value units. Error bars represent an average standard deviation of 2.5%, as determined by repetitive measurements of individual bands (instrument error). Each graph is representative of at least two separate experiments.

In contrast to results with 21PT and 21NT cells, anti- $\alpha\beta 5$ and $\beta 1$ integrin antibodies showed no blocking effect on OPN-induced cell migration of MDA-MB-435 cells, when used at the same high concentrations shown to effect complete blocking of 21PT and 21NT responsiveness (15 $\mu\text{g}/\text{ml}$ of either anti-integrin antibody; Fig. 2). On the other hand, OPN-induced migration of MDA-MB-435 cells was completely blocked by anti- $\alpha\beta 3$ integrin antibody, at a concentration (25 $\mu\text{g}/\text{ml}$) lower than that which still had no effect on migration of 21PT or 21NT (30 $\mu\text{g}/\text{ml}$; cf. Fig. 1; $P = 0.0008$, Student's *t*-test).

The OPN-induced migration of the metastatic cell line of this series—MDA-MB-435, thus was found to be $\alpha\beta 3$ integrin-dependent, whereas that of non-metastatic 21NT and 21PT cells was $\alpha\beta 5$ and $\beta 1$ -dependent, $\alpha\beta 3$ -independent.

HGF-Induced Cell Migration and Synergistic Effect With OPN

As was found for response to OPN, all three cell lines (21PT, 21NT, MDA-MB-435) showed increased cell migration in response to human recombinant HGF alone (Fig. 3). Combining both HGF and osteopontin in the lower chamber resulted in a degree of cell migration for all three cell lines that was significantly greater than the sum of the isolated HGF and OPN responses (i.e., synergistic; Fig. 3; $P < 0.02$ for all, Student's *t*-test).

Induction of Met (HGFR) Activity by HGF and OPN

Treatment of 21PT and 21NT cells with either HGF or OPN (Fig. 4a,b) resulted in rapid activation of total Met kinase activity in both instances (after 20 min of HGF stimulation, 30 min of OPN stimulation). For both 21PT and 21NT cells, the increase in total Met kinase activity with OPN treatment (at 30 min) was followed by an increase in Met protein level at 5 h of incubation with OPN (Fig. 4c).

Treatment of MDA-MB-435 cells with HGF or OPN (Fig. 5a) resulted in an increase in specific Met kinase activity after 20 (for HGF) to 30 (for OPN) min of incubation, which was associated with an increased tyrosine phosphorylation of Met as well (Fig. 5b). In contrast with 21PT and 21NT cells, we have not been able to detect an OPN-induced increase in total




Fig. 5. Induction of specific Met kinase activity (a) and Met tyrosine phosphorylation (b) of MDA-MB-435 cells by HGF and OPN. MDA-MB-435 cells were incubated $\pm 50 \mu\text{g}/\text{ml}$ hrOPN or 20 ng/ml HGF for the times indicated. Cell lysates were prepared, and equal protein amounts of each lysate were immunoprecipitated with anti-Met IgG. As levels of total Met protein were higher and fluctuated more in MDA-MB-435 cells than in 21PT and 21NT, activation of Met protein in MDA-MB-435 was more appropriately expressed as Relative Met kinase activity (a) and tyrosine phosphorylation (b). a: In vitro Met kinase activity was assayed as described in Materials and Methods. Relative Met kinase activity, normalized to total Met protein, was quantitated using a Phosphorimager. b: Immunoprecipitates were subjected to 7% SDS-PAGE and transferred to nitrocellulose. The membrane was blocked with 1% BSA in TBST, and probed with anti-phosphotyrosine antibody. Detection was performed with HRP-labeled anti-mouse antibody and ECL. Relative Met tyrosine-phosphorylation normalized to total Met protein was quantitated by densitometry. Error bars represent an average standard deviation of 5.0%, as determined by repetitive measurements of individual bands (instrument error). Each graph is representative of at least two separate experiments.

Integrin-Dependent, OPN-Induced Migration Involves Met

471




Fig. 6. Time course showing effect of OPN on expression of HGF and Met mRNA by 21PT (PT), 21NT (NT), and MDA-MB-435 (MDA) cells. Near-confluent (85–90%) cell cultures were incubated in serum-free medium with 50 µg/ml OPN for 0, 5, 18, or 24 h. Total RNA (10 µg/lane) was analysed by Northern blotting for expression of HGF (6.0 kb) or Met (HGFR; 8.0 kb [full-length transcript, upper band], 7.0 kb [lower band]). RNA loading and integrity were verified by assessment of 18S rRNA (2.1 kb). Level of HGF and Met (both 8kb and 7kb transcripts) are shown in relation to 18S rRNA, expressed as the ratio of densitometry values for the respective bands (HGF:18S, Met:18S).

Met protein levels in MDA-MB-435 cells, although basal level of Met protein expression in MDA-MB-435 is higher than in 21PT or 21NT (data not shown).

Time Course Showing Effect of OPN on Expression of HGF and Met mRNA

21PT, 21NT, and MDA-MB-435 cells treated with OPN (50 µg/ml) for 0–24 h (Fig. 6), all showed low basal levels of HGF mRNA. Only slight increase in HGF mRNA was detected for 21PT and 21NT, with no appreciable increase for MDA-MB-435 cells (by 18–24 h). In contrast, levels of Met RNA were found to significantly increase in all three cell lines between 5 and 18 h of OPN exposure, falling off by 24 h. Thus, although little change in HGF mRNA was seen after up to 24 h of OPN exposure, significant induction of Met mRNA was seen for all three cell lines.

DISCUSSION

OPN has been implicated in the malignancy of breast cancer in a number of recent studies [e.g., Oates et al., 1996; Singhal et al., 1997; Sung et al., 1998; Tuck et al., 1998, 1999]. It has been shown to be involved in cell adhesion of MECs, and can also induce cell migration in an RGD-dependent fashion [Xuan et al., 1994, 1995; Senger et al., 1996]. The HGF/Met path-

way has also been associated with breast cancer malignancy [Yamashita et al., 1994; Tuck et al., 1996; Yao et al., 1996; Jin et al., 1997; Beviglia et al., 1997; Ghoussoub et al., 1998], and is a potent inducer of MEC motility [Bhargava et al., 1992; Rosen et al., 1994; Rahimi et al., 1998]. Here we examine the nature of the integrin response to OPN, in order to establish the specific cell surface integrins involved. We also show that OPN-induced cell migration involves activation of the HGF receptor in a synergistic fashion with HGF, consistent with cross-talk between integrin and growth factor mediated pathways.

Our discovery that, in a series of breast epithelial cells of differing degrees of malignancy, different cell surface integrins may couple with Met in inducing cell migration is a novel finding. The metastatic member of the series studied, MDA-MB-435 cells, showed the most marked synergy between OPN and HGF in the migration response, and migrated in an $\alpha v \beta 3$, not $\alpha v \beta 5$ or $\beta 1$ -dependent fashion. In contrast, the non-metastatic cell lines, 21PT and 21NT, migrated in an $\alpha v \beta 5$ and $\beta 1$ dependent, $\alpha v \beta 3$ -independent fashion. In support of this finding is the work of Wong et al. [1998], who reported that MDA-MB-435 cells express $\alpha v \beta 3$ integrin, while less malignant MDA-MB-231 and MCF-7 cells do not (although they all express $\alpha v \beta 5$ and

$\beta 1$). Similarly, van der Pluijm et al. [1997] reported higher expression of $\alpha v \beta 3$ in more malignant members of a series of breast carcinoma cell lines. A specific association between $\alpha v \beta 3$ expression and breast cancer metastasis has also been reported by Liapis et al. [1996], who detected $\alpha v \beta 3$ integrin expression in 100% of breast carcinomas that had metastasized to bone.

This difference in integrin utilization of cells at different stages of progression could affect malignancy in a number of different ways. For example, $\alpha v \beta 3$ may be necessary for specific adhesion events vital to invasion and metastasis; a breast cancer cell initially expressing $\alpha v \beta 1$ or $\alpha v \beta 5$ may require activation of $\beta 3$ in order to complete that step of the metastatic cascade. Evidence in favor of this scenario (at least in the case of melanoma) comes from the work of Nip et al. [1992], who showed that binding of metastatic cells to lymph node matrix depends on $\alpha v \beta 3$ interactions. Alternatively, different integrins may be coupled to different signal transduction pathways, with $\alpha v \beta 3$ specifically required for activation of a particular set of genes important in aspects of invasion and metastasis. Ligation of $\alpha v \beta 3$ for example, has been shown to induce MMP-2 expression and invasion of melanoma cells [Seftor et al., 1992; Bafetti et al., 1998]. Whether or not different integrins couple to Shc can influence activation of transcription from the Fos serum response element (SRE), affecting responsiveness to growth factors [Wary et al., 1996]. In the case of coupling with the HGF/Met pathway, our work suggests that OPN-induced cell migration via either non- $\alpha v \beta 3$ integrins (21PT, 21NT) or $\alpha v \beta 3$ integrin (MDA-MB-435) is associated with Met activation, but that the synergistic effect on cell migration is much more pronounced in the cells (MDA-MB-435) expressing $\alpha v \beta 3$.

We also found that OPN-induced migration of all three cell lines involves activation of the HGF receptor, Met, with an initial increase in Met activity followed by an increase in Met RNA expression. The kinetics of this effect differ slightly for MDA-MB-435 vs. the 21T series cells. For 21PT and 21NT, a detectable increase in Met protein was also found. Although a similar increase in Met protein levels of MDA-MB-435 cells did not occur, the basal level of Met protein in these

cells is quite high, and as specific activity of Met is substantially increased with OPN induction, it is possible that Met turnover is such that protein levels do not further accumulate as they do for the 21T series cells. Regardless, Met is activated by OPN in all three cell lines, and this is associated with increased cell migration. In contrast, HGF mRNA levels were low in all three cell lines, and showed little or no change with OPN treatment. Furthermore, we have not detected increased HGF activity in conditioned media of OPN-treated 21T series or MDA-MB-435 cells (data not shown). Activation of Met by OPN is thus likely due to either an increased sensitivity to trace amounts of ligand present, or to ligand-independent activation. Ligand-independent activation of Met by cellular adhesion has been previously reported for melanoma cells, although the cell surface adhesion receptors involved were not examined [Wang et al., 1996]. Furthermore, integrin binding has been shown to be essential for growth factor (EGF, PDGF, bFGF, IGF-1) induced signal transduction and cell migration [Miyamoto et al., 1996; Brooks et al., 1997]. Reciprocally, HGF can activate cell surface integrins and hence cellular adhesion (and motility) [van der Voort et al., 1997; Weimar et al., 1997, 1998; Trusolino et al., 1998]. Thus, a two-way interaction between integrin and growth factor-mediated pathways likely occurs in the induction of cellular responses such as cell migration.

Multiple points of interaction between signal transduction pathways activated by integrins vs. growth factors have been identified [reviewed in Sastry and Horowitz, 1996; Giancotti, 1997; Swartz, 1997]. Cell attachment can enhance autophosphorylation of growth factor receptors (EGFR, PDGFR, and now Met) in response to their cognate ligands. Integrin binding also has been found to activate phospholipase C (and hence protein kinase C), Raf, and/or MEK in the MAP kinase pathway, and PI-3 kinase in the PI-3K/Rac pathway. All of these pathways are also influenced by growth factors, although the synergistic relationship reported here would suggest that growth factor receptors and integrins may act at different points in the pathway. For example, it has been shown that fibronectin binding to cell surface integrin activates synthesis and supply of phosphatidylinositol 4,5 biphosphate, whereas PDGF receptor controls the activity of phospho-

lipase C [McNamee et al., 1992]. The physical association of integrins and growth factor receptors at the focal adhesion complex (FAC) [Plopper et al., 1995] provides a mechanism by which such cross-talk would be facilitated. Finally, interactions between growth factor and integrin pathways could occur at the level of differential effects on members of the FAC itself [Schlaepfer and Hunter, 1998].

In the case of signal transduction initiated by OPN, ligation of $\alpha v \beta 3$ by OPN activates PI-3 kinase in osteoclasts [Hruska et al., 1995]. In an osteoblastic cell line (UMR 106-8), OPN triggers the autophosphorylation of focal adhesion kinase (FAK) [Liu et al., 1997]. In *ras*-transformed fibroblasts, OPN can induce tyrosine phosphorylation of a number of different FAC associated proteins [Lopez et al., 1995]. HGF activation of its receptor, Met, can also stimulate phosphorylation of FAK in some cells, perhaps via $pp60^{src}$ [Chen et al., 1998]. We have shown in this report that OPN is also capable of activating Met. Thus, although our understanding of the signal transduction pathways induced by OPN is yet in early stages, multiple points of potential interaction between the integrin and growth factor receptor pathways involved are already beginning to emerge.

The interactions between different integrin pathways induced by OPN and the HGF/Met growth factor pathway not only helps conceptually in understanding the clinical associations we have observed between OPN, HGF and malignancy, but also provides clues to regulatory processes vital to tumor aggressiveness—prime targets for treatment strategies based on blocking these processes.

ACKNOWLEDGMENTS

Dr. Tuck is supported by a "Career Development" Award from the U.S. Army Breast Cancer Research Program.

REFERENCES

- Assiian RK. 1997. Anchorage-dependent cell cycle progression. *J Cell Biol* 136:1-4.
- Bafetti LM, Young TN, Itoh Y, Stack MS. 1998. Intact vitronectin induces matrix metalloproteinase-2 and tissue inhibitor of metalloproteinases-2 expression and enhanced cellular invasion by melanoma cells. *J Biol Chem* 273:143-149.
- Band V, Zajchowski D, Swisshelm K, Trask D, Kulesa V, Cohen C, Connolly J, Sager R. 1990. Tumor progression in four mammary epithelial cell lines derived from the same patient. *Cancer Res* 50:7351-7357.
- Beviglia L, Matsumoto K, Lin CS, Ziober BL, Kramer RH. 1997. Expression of the α -Met/HGF receptor in human breast carcinoma: Correlation with tumor progression. *Int. J. Cancer* 74:301-309.
- Bhargava M, Joseph A, Knesal J, Halaban R, Li Y, Pang S, Goldberg I, Setter E, Donovan MA, Zarnegar R, Michalopoulos GA, Nakamura T, Faletto D, Rosen EM. 1992. Scatter factor and hepatocyte growth factor: Activities, properties, and mechanism. *Cell Growth Differ* 3:11-20.
- Birchmeier C, Gherardi E. 1998. Developmental roles of HGF/SF and its receptor, the α -Met tyrosine kinase. *Trends Cell Biol* 8:404-410.
- Bissell MJ, Weaver VM, Lelievre SA, Wang F, Petersen OW, Schmeichel KL. 1999. Tissue structure, nuclear organization, and gene expression in normal and malignant breast. *Cancer Res* 59:1757-1763.
- Bourguignon LY, Gunja-Smith Z, Iida N, Zhu HB, Young LJ, Muller WJ, Cardiff RD. 1998. CD44v(3,8-10) is involved in cytoskeleton-mediated tumor cell migration and matrix metalloproteinase (MMP-9) association in metastatic breast cancer cells. *J Cell Physiol* 176:206-215.
- Bourguignon LY, Zhu H, Shao L, Zhu D, Chen YW. 1999. Rho-kinase (ROK) promotes CD44v(3,8-10)-ankyrin interaction and tumor cell migration in metastatic breast cancer cells. *Cell Motil Cytoskeleton* 43:269-287.
- Brooks PC, Klempke RL, Schön S, Lewis JM, Schwartz MA, Cheresah DA. 1997. Insulin-like growth factor receptor cooperates with integrin $\alpha v \beta 5$ to promote tumor cell dissemination in vivo. *J Clin Invest* 99:1890-1899.
- Chen HC, Chan PC, Tang MJ, Chang CH, Chang TJ. 1988. Tyrosine phosphorylation of focal adhesion kinase stimulated by hepatocyte growth factor leads to mitogen-activated protein kinase activation. *J Biol Chem* 273:25777-25782.
- Ghoussoub RA, Dillon DA, D'Aquila T, Rimm EB, Fearon ER, Rimm DL. 1998. Expression of α -met is a strong independent prognostic factor in breast carcinoma. *Cancer* 82:1519-1520.
- Giancotti FG. 1997. Integrin signaling: Specificity and control of cell survival and cell cycle progression. *Curr Opin Cell Biol* 9:691-700.
- Gumbiner BM. 1996. Cell adhesion: the molecular basis of tissue architecture and morphogenesis. *Cell* 84:345-357.
- Heldin CH, Ostman A, Ronnstrand L. 1998. Signal transduction via platelet-derived growth factor receptors. *Biochim Biophys Acta* 1378:F79-113.
- Hruska K, Rolnick F, Huskey M, Alvarez U, Cheresah D. 1996. Engagement of the osteoclast integrin $\alpha v \beta 3$ by osteopontin stimulates phosphatidylinositol 3-hydroxyl kinase activity. In: Denhardt DT, Butler WT, Chambers AF, Senger DR, editors. Osteopontin—role in cell signaling and adhesion. *Ann NY Acad Sci* 760:151-165.
- Hu DD, Lin EC, Kovach NL, Hoyer JR, Smith JW. 1995. A biochemical characterization of the binding of osteopontin to integrins $\alpha v \beta 1$ and $\alpha v \beta 5$. *J Biol Chem* 270:26232-26238.
- Jin L, Fuchs A, Schnitt SJ, Yao Y, Joseph A, Lamszus K, Park M, Goldberg ID, Rosen EM. 1997. Expression of scatter factor and α -met receptor in benign and malignant breast tissue. *Cancer* 79:749-760.

- Katagiri YU, Sleeman J, Fujii H, Harrlich P, Hotta H, Tanaka K, Chikuma S, Yagita H, Okumura K, Murakami M, Said I, Chambers AF, Ueda T. 1999. CD44 variants but not CD44s cooperate with beta1-containing integrins to permit cells to bind to osteopontin independently of arginine-glycine-aspartic acid, thereby stimulating cell motility and chemotaxis. *Cancer Res* 59:219-228.
- Lisapis H, Flath A, Kitazawa S. 1996. Integrin $\alpha v \beta 3$ expression by bone-residing breast cancer metastases. *Diag Mol Pathol* 5:127-135.
- Liaw L, Skinner MP, Raines EW, Ross R, Cheresch DA, Schwartz SM, Giachelli CM. 1995. The adhesive and migratory effects of osteopontin are mediated via distinct cell surface integrins. Role of alpha v beta 3 in smooth muscle cell migration to osteopontin in vitro. *J Clin Invest* 95:713-724.
- Liu Y-K, Uamara T, Nemoto A, Yabe T, Fuhii N, Ushida T, Tateishi T. 1997. Osteopontin involvement in integrin-mediated cell signaling and regulation of expression of alkaline phosphatase during early differentiation of UMR cells. *FEBS Lett* 420:112-116.
- Lopez CA, Davis RL, Mou K, Denhardt DT. 1996. Activation of a signal transduction pathway by osteopontin. In: Denhardt DT, Butler WT, Chambers AF, Senger DR, editors: *Osteopontin—role in cell signaling and adhesion*. Ann NY Acad Sci 760:324-326.
- McNamee HM, Ingber DE, Schwartz MA. 1992. Adhesion to fibronectin stimulates inositol lipid synthesis and enhances PDGF-induced inositol breakdown. *J Cell Biol* 121:673-678.
- Miyamoto S, Teramoto H, Gutkind JS, Yamada KM. 1996. Integrins can collaborate with growth factors for phosphorylation of receptor tyrosine kinases and MAP kinase activation: Roles of integrin aggregation and occupancy of receptors. *J Cell Biol* 135:1633-1642.
- Nakamura T, Nishizawa T, Hagiya M, Seki T, Shimonishi M, Sugimura A, Tashiro K, Shimizu S. 1989. Molecular cloning and expression of hepatocyte growth factor. *Nature* 34:440-443.
- Nip J, Shibata H, Laskutoff DJ, Cheresch DA, Brodt P. 1992. Human melanoma cells derived from lymphatic metastases use integrin alpha v beta 3 to adhere to lymph node vitronectin. *J Clin Invest* 90:1406-1413.
- Oates AJ, Barraclough R, Rudland PS. 1996. The identification of osteopontin as a metastasis-related gene product in a rodent mammary tumour model. *Oncogene* 13: 97-104.
- Park M, Dean M, Kaul K, Braun MJ, Gonda MA, Vande Woude G. 1987. Sequence of MET protooncogene cDNA has features characteristic of the tyrosine kinase family of growth factor receptors. *Proc Natl Acad Sci USA* 84: 6879-6883.
- Plopper G, McNamee HP, Dike LE, Bojanowski K, Ingber DE. 1995. Convergence of integrin and growth factor receptor signaling pathways within the focal adhesion complex. *Mol Biol Cell* 6:1349-1365.
- Price JE, Polyzos A, Zhang RD, Daniels LM. 1990. Tumorigenicity and metastasis of human breast carcinoma cell lines in nude mice. *Cancer Res* 50:717-721.
- Rahimi N, Hung W, Tremblay E, Saulnier R, Elliott B. 1998. c-Src kinase activity is required for hepatocyte growth factor-induced motility and anchorage-independent growth of mammary carcinoma cells. *J Biol Chem* 273:33714-33721.
- Rosen EM, Knessel J, Goldberg ID, Jin L, Bhargava M, Joseph A, Zitnik R, Wines J, Kelley M, Rockwell S. 1994. Scatter factor modulates the metastatic phenotype of the KMT6 mouse mammary tumor. *Int J Cancer* 57:706-714.
- Sastry SK, Horowitz AF. 1998. Adhesion - growth factor interactions during differentiation: An integrated biological response. *Dev Biol* 180:455-467.
- Schlaepfer DD, Hunter T. 1998. Integrin signalling and tyrosine phosphorylation: Just the FAKs? *Trends Cell Biol* 8:151-157.
- Schwartz MA. 1997. Integrins, oncogenes, and anchorage independence. *J Cell Biol* 139:575-578.
- Seedorf K. 1995. Intracellular signaling by growth factors. *Metabolism* 44:24-32.
- Seftor RE, Seftor EA, Gehlsen KR, Stetler-Stevenson WG, Brown, PD, Ruoslahti E, Hendrix MJ. 1992. Role of the alpha v beta 3 integrin in human melanoma cell invasion. *Proc Natl Acad Sci* 89:1557-1561.
- Senger DR, Ferruzzi CA. 1996. Cell migration promoted by a potent GRGDS-containing thrombin-cleavage fragment of osteopontin. *Biochim Biophys Acta* 1314:13-24.
- Singhal H, Bantiata DS, Tonkin KS, O'Malley FP, Tuck AB, Chambers AF, Harris JP. 1997. Elevated plasma osteopontin in metastatic breast cancer associated with increased tumor burden and decreased survival. *Clin Cancer Res* 3:605-611.
- Sung V, Gillis C, Murray A, Clarke R, Aaron AD, Azumi N, Thompson EW. 1998. The LCC15-MB human breast cancer cell line expresses osteopontin and exhibits an invasive and metastatic phenotype. *Exp Cell Res* 241:273-284.
- Trusolino L, Sarini G, Cecchini G, Beanti C, Ambesi-Impombato FS, Marchisio PC, de Filippi R. 1998. Growth factor-dependent activation of $\alpha v \beta 3$ integrin in normal epithelial cells: Implications for tumor invasion. *J Cell Biol* 142:1145-1156.
- Tuck AB, Park M, Sterns EE, Buzg A, Elliott BE. 1996. Coexpression of hepatocyte growth factor and receptor (met) in human breast carcinoma. *Am J Pathol* 148:225-232.
- Tuck AB, O'Malley FP, Singhal H, Harris JP, Tonkin KS, Kerkvliet N, Saad Z, Doig GS, Chambers AF. 1998. Osteopontin expression in a group of lymph node negative breast cancer patients. *Int J Cancer* 79:502-508.
- Tuck AB, Arseneault DM, O'Malley FP, Hota C, Ling MC, Wilson SM, Chambers AF. 1999. Osteopontin induces increased invasiveness and plasminogen activator expression of human mammary epithelial cells. *Oncogene* 18:4237-4246.
- van der Pluijm G, Vloedgraven H, Papapoulos S, L6wik C, Grzesik W, Kerr J, Robey PG. 1997. Attachment characteristics and involvement of integrins in adhesion of breast cancer cell lines to extracellular bone matrix components. *Lab Invest* 77:665-678.
- van der Voort R, Tohor TEI, Keenen RMJ, Smit L, Groenink M, Pals ST. 1997. Paracrine regulation of germinal center B cell adhesion through the c-Met-hepatocyte growth factor/scatter factor pathway. *J Exp Med* 185: 2121-2131.

Integrin-Dependent, OPN-Induced Migration Involves Met

475

- Vande Woude GF, Jeffers M, Cortner J, Alvord G, Tsarfaty I, Rouseau J. 1997. Met-HGF/SF: Tumorigenesis, invasion and metastasis. *Ciba Found Symp* 212:148-154.
- Wang R, Kobayashi R, Bishop JM. 1996. Cellular adherence elicits ligand-independent activation of the Met cell-surface receptor. *Proc Natl Acad Sci*. 93:8425-8430.
- Wary KK, Mainiero F, Isakoff SJ, Marcantonio EE, Giannotti FG. 1996. The adaptor protein Shc couples a class of integrins to the control of cell cycle progression. *Cell* 87:733-743.
- Weber GF, Ashkar S, Glimcher MJ, Cantor H. 1996. Receptor-ligand interaction between CD44 and osteopontin (Eta-1). *Science* 271:509-512.
- Weimar IS, de Jong D, Muller EJ, Nakamura T, Gorp JMHH, de Gast GC, Gerritsen WR. 1997. Hepatocyte growth factor/scatter factor promotes adhesion of lymphoma cells to extracellular matrix molecules via $\alpha 4 \beta 1$ and $\alpha 5 \beta 1$ integrins. *Blood* 89:990-1000.
- Weimar IS, Miranda N, Muller EJ, Hakman A, Kerst JM, de Gast GC, Gerritsen WR. 1998. Hepatocyte growth factor/scatter factor (HGF/SF) is produced by human bone marrow stromal cells and promotes proliferation, adhesion and survival of human hematopoietic progenitor cells (CD34+). *Exp Hematol* 26:885-894.
- Wong NC, Mueller BM, Barbas CF, Ruminaki P, Quaranta V, Lin ECK, Smith JW. 1998. αv integrins mediate adhesion and migration of breast carcinoma cell lines. *Clin Exp Metastasis* 16:50-61.
- Xuan JW, Hota C, Chambers AF. 1994. Recombinant GST-human osteopontin fusion protein is functional in RGD-dependent cell adhesion. *J Cell Biochem* 54:247-255.
- Xuan JW, Hota C, Shigeyama Y, D'Errico JA, Somerman MJ, Chambers AF. 1995. Site-directed mutagenesis of the arginine-glycine-aspartic acid sequence in osteopontin destroys cell adhesion and migration functions. *J Cell Biochem* 57:680-690.
- Yamashita J, Ogawa M, Yamashita S, Nomura K, Kuramoto M, Saishoji T, Shin S. 1994. Immunoreactive hepatocyte growth factor is a strong and independent predictor of recurrence and survival in human breast cancer. *Cancer Res* 54:1630-1633.
- Yao Y, Jin L, Fuchs A, Joseph A, Hastings HM, Goldberg ID, Rosen EM. 1996. Scatter factor protein levels in human breast cancers: Clinicopathological and biological correlations. *Am J Pathol* 149:1707-1717.